



Scientific article



## GC-MS Evaluation of stem bark extract of *sterculia oblonga* for biomedical potentials

Patience Yake Nkom <sup>a</sup>, Esther Offiong Asuquo <sup>a</sup>, Lynius Ashipu Bebiaundeye <sup>a</sup> and Endurance Egwu Ukam <sup>a</sup>

<sup>a</sup> *University of Cross River State, Calabar, Nigeria*

### ITEM INFORMATION

#### Item history:

Received on May 13, 2024  
Accepted on June 26, 2024

#### Keywords:

Bioactive compounds  
*Sterculia oblonga*  
Organo-sulphur  
Phenols  
Hydrocarbons

### ABSTRACT

The stem-bark of *sterculia oblonga*, which is a rich lignocellulosic material, was analysed using GC-MS chromatography in order to identify the bioactive compounds present in the plant and deduce its possible therapeutic potentials. The pulverized plant sample was extracted by maceration for 48 hours using ethanol solvent, after solvent recovery, the crude extract was used for the GC-MS analysis. Results revealed the presence of 12 bioactive compounds namely: 1-(Butylsulfanyl)-2-ethoxyethane 4-methoxyphenol (mequinol), 2,6-dimethoxy-phenol (Syringol), 3,4,5-Trimethoxyphenol, 4,4,5,6-Tetramethyltetrahydro-1,3-oxazin-2-thione, n-Hexadecanoic acid (Palmitic acid), 1-Docosene, 14-methyl-8-hexadecen-1-ol, 1,19-Eicosadiene, Z-10-Tetradecen-1-ol acetate, 1-Nonadecene and Oleic Acid. These chemicals were further grouped into 5 viz organo-sulphur (2%), phenols (24%), alkaloids (2%), fatty acids/esters (22.18%) and hydrocarbons (8.66%). Because of the high percentage of phenolic compounds present, the ethanolic extract of stem-bark of *Sterculia oblonga* could find useful therapeutic applications as antimicrobial, anti-cancer, anti-oxidant and anti-diabetic agents.

© 2024 Professionals On Line sac. Perú World Health Journal

All rights reserved



<https://doi.org/10.47422/whj.v5i1.46>



## INTRODUCCIÓN

The biopolymers called hemicellulose, cellulose and lignin, which have demonstrated antimicrobial properties, make up most lignocellulosic materials<sup>1</sup>. One of the most prevalent natural materials, lignocellulosic materials have favorable qualities like low density and biodegradability. They are also readily available, cheap, and environmentally friendly. Massive amounts of lignocellulose are extracted from agricultural waste, but only a small amount of that is used since lignocellulose's biotechnological significance is not well understood<sup>2</sup>. The amount of lignocellulosic content available in a material varies. Considering the enormous amount of chemical compounds available in the extract from these biopolymeric materials, there are countless prospects for the development of new drugs<sup>3</sup>. For instance, example aqueous extracts from willow leaves (*Salix sasaf*) have been reported to be effective against Guman carcinoma cells and have been tested in vivo and in vitro<sup>4</sup>; antimicrobial activity of *Punica granatum* has been reported to be effective against all test microorganisms<sup>5</sup>. The heavy dependency of people on traditional medicine for primary health care has prompted in-depth research into plants to enhance their curative properties<sup>6</sup>. Some common sources of lignocellulosic materials are shown in Fig. 1.



Fig. 1: common sources of lignocellulosic materials

Yellow sterculia, or *Sterculia oblonga* (Fig. 2), is a commercial wood tree in the Malvaceae family that is found mostly in tropical rainforests in countries like Cameroon, Gabon, Liberia, Ghana, Nigeria, etc. It is reported that the edible seeds have a pea-like flavor. They are best

eaten roasted or raw. Handbags, hats, rope, paper, and other items are made from the bark's fibers. The wood can be used to build little furniture pieces, like cupboards, or matchsticks.



Fig 2: pictural view of *Sterculia oblonga*

The dried wood of *Sterculia oblonga* has been reported to contain 11.5 – 12.4% furfural; 36.2 – 46.32 cellulose; 19.8-21.2% pentosans; 18.2-22.7% lignin; and 1.3-3.2% ash<sup>7</sup>. Ezeonu and Ejikeme (2016)<sup>8</sup> also reported that *sterculia oblonga* contain tannins, 124 mg/100g; flavonoids, 3.4%; alkaloids, 10%; cyanide, 1.66 mg/100g; lipids 4.8%; phenol, 620 mg/g; and oxalate, 1.31 g/100g. The gas chromatography-mass spectroscopy analysis of essential oil from *sterculia oblonga* shows that the plant contains 33 compounds ranging from simple compound like p-xylene to very complex compounds like diisocetylphthalate<sup>9</sup>. Report on petroleum spirit extract showed that the plant contains 25 natural products which include 2-myristgnoyl pantetheine; D-limonene, limonene-b-0l, pivalate, chalestan-3-0l, 2-methylene-, (3 $\beta$ , 5 $\alpha$ ), cis-13-eicosenoic acid, Hexadecane, I, 1-bis(dedecyloxy), Z-plytol, and 17-pentatriacontene<sup>10</sup>. This study aimed to analyze the ethanolic extract of *Sterculia oblonga* stem bark via gas chromatography-mass spectrometry (GC-MS) to determine their probable usage in pharmacology and ascertain their phytochemical composition.

## METHODOLOGY

**Plant sample:** The fresh stem bark of yellow sterculia (*Sterculia oblonga*) was collected from the rain forest of Akamkpa Local Government Area, Cross River State, Nigeria. The identification and authentication of the plant sample was done by the Herbarium Unit of Botany Department, University of Calabar. It was washed clean with distilled water, air-dried, crushed into tiny pieces, and stored at room temperature in air-tight containers for further studies.

**Extraction:** The maceration method of extraction was used in this study because of its easy method, no heat involved, suitable for thermally sensitive materials, inexpensive installation and maintenance costs. A

total of 850 g of the dried plant was extracted sequentially using ethanol by cold maceration for 48 hours. The resulting solution was filtered and concentrated to dryness using rotary evaporator.

Determination of the secondary metabolites of the crude extract using gas chromatography-mass spectrometry (GC-MS): The identification of the individual compounds of the ethanol crude extract was done by standard method using a GC-MS analyzer. The components were identified by correlating the masses of the generated spectra to those of the library's reference mass spectra.

## RESULTS AND DISCUSSION

Results in Table 1 show the yield of the three extracts and the calculated percentage yield for the three extracts. Gas chromatogram of ethanol extract of *Sterculia oblonga* stem bark is presented in Fig. 3. The GC-MS analysis of bioactive compounds in ethanol extract of *sterculia oblonga* stem bark and pharmacological potentials of the phyto-compounds in the ethanolic extract of *sterculia oblonga* stem bark are presented in Table 2 and 3 respectively.

**Table 1**

Percentage yields of extract

Solvent	Weight of plants (g)	Weight of extract (g)	Colour of extract	Texture of extract	Percentage yield (%)
Ethanol	1200	13.21	Green	Sticky	11




Fig. 3: Gas chromatogram of ethanol extract of *Sterculia oblonga* stem bark

**Table 2**

GC-MS results of Secondary metabolites in Ethanol Ethanol Extract of *Sterculia Oblonga* Stem Bark

S/N	Retention Time	Peak area (%)	Molecular formula	MW	Name of compound
1.	3.37	1.00	C <sub>8</sub> H <sub>18</sub> OS	162	1-(Butylsulfanyl)-2-ethoxyethane
2.	4.60	1.56	CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub> OH	124	4-methoxyphenol (mequinol)
3.	7.72	0.90	HO(CH <sub>3</sub> O) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	154	2,6-dimethoxy-phenol (Syringol)
4.	10.48	1.12	C <sub>9</sub> H <sub>12</sub> O <sub>4</sub>	184	3,4,5-Trimethoxyphenol
5.	12.95	0.93	C <sub>8</sub> H <sub>15</sub> NOS	173	4,4,5,6-Tetramethyltetrahydro-1,3-oxazin-2-thione
6.	13.55	9.98	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256	n-Hexadecanoic acid (Palmitic acid)
7.	16.74	1.14	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	308	1-Docosene
8.	17.92	2.75	C <sub>17</sub> H <sub>34</sub> O	254	14-methyl-8-hexadecen-1-ol
9.	18.25	1.53	C <sub>20</sub> H <sub>38</sub>	278	1,19-Eicosadiene
10.	19.99	8.80	C <sub>16</sub> H <sub>30</sub> O <sub>2</sub>	254	Z-10-Tetradecen-1-ol acetate
11.	20.19	3.24	C <sub>19</sub> H <sub>38</sub>	266	1-Nonadecene
12.	20.64	3.40	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282	Oleic Acid

**Table 3**

*Pharmacological potentials of the Phyto-compounds in the Ethanolic Extract of Sterculia Oblonga Stem Bark*

S/N	Name of compound	Class of phytochemical	Major function	Pharmacological activity
	1-(Butylsulfanyl)-2-ethoxyethane	Organo-sulfur	Antibiotic and antidiabetic agent	Anti-cancer agent <sup>11</sup> , fertility activity <sup>27</sup>
	4-methoxyphenol (mequinol)	Phenol	Antimicrobial	Diabetic Wound dressing <sup>12</sup> , cosmetic preservative <sup>13</sup>
	2,6-dimethoxy-phenol (Syringol)	Phenol	Antimicrobial	Antitermitic activity <sup>14</sup>
	3,4,5-Trimethoxyphenol	Phenol	Antimicrobial	Anti-invasive cancer agents <sup>15</sup> , Antiplasmodial activity <sup>16</sup>
	4,4,5,6-Tetramethyltetrahydro-1,3-oxazin-2-thione	Alkaloid	Antioxidant, Antimicrobial	Cosmetics, anti-cancer and anti-cholesterol activity <sup>17</sup>
	n-Hexadecanoic acid (Palmitic acid)	Fatty acids/ Ester	Anti-inflammation	Anti-inflammation <sup>18</sup> , Cytotoxic activity <sup>19</sup>
	1-Docosene	Hydrocarbon	Antimicrobial	Antibacterial <sup>20</sup> , antioxidant <sup>21</sup> and antifungal activity <sup>22</sup>
	14-methyl-8-hexadecen-1-ol	Hydrocarbon	-	No therapeutic report
	1,19-Eicosadiene	Hydrocarbon	Antimicrobial	Antidiabetic activity
	Z-10-Tetradecen-1-ol acetate	Ester	sex pheromone	Sex attractant <sup>23</sup>
	1-Nonadecene	Hydrocarbon	Anti-inflammatory	Antimicrobial, antioxidant, anticancer <sup>24</sup>
	Oleic Acid	Fatty acid	Antimicrobial; Anti-inflammatory	Anti-inflammation, modulation of leukocytes activity <sup>25,26</sup>




Fig. 4: Percentage composition of phytochemicals identified in the crude Ethanolic extract of Sterculia Oblonga Stem Bark

The GC-MS analysis of the stem-bark of *Sterculia oblonga* showed the presence of twelve (12) different phytochemicals including organo-Sulphur compounds (2%), phenols (24%), alkaloids (2%), fatty acids/esters (52%) and hydrocarbons (20%) (Fig. 4). From the results, it is safe to say that the stem-bark of *Sterculia oblonga* contains very high amount of fatty acids/esters, followed by phenols, and

hydrocarbons. The presence of these compounds sums up the effective use of the plant in many pharmacological applications such as wound healing, anti-cancer, anti-microbial, anti-oxidant and sex stimulant activities.



Organosulfur compounds have shown antimicrobial activity against various pathogens, including bacteria, viruses, fungi, and parasites. These compounds can help inhibit the growth of harmful microorganisms and may be beneficial in fighting infections<sup>28</sup>. A study by Kamenetsky *et al.* (2005)<sup>27</sup> revealed that organosulfur compounds could have fertility potential<sup>27,29</sup>. Other therapeutic applications of Organosulfur compounds include: Anti-cancer<sup>11</sup>, antioxidant activity<sup>30</sup>. However, the GC-MS chromatogram of the stem-bark of *Sterculia oblonga* revealed only 2% of organo-sulfur content.

The presence of phenolic compounds in medicinal plants contributes significantly to their therapeutic potential. These compounds exhibit antioxidant, anti-inflammatory, antimicrobial, and other bioactive properties that can help prevent or alleviate various health conditions. By consuming plant-based foods rich in phenolic compounds or using herbal remedies derived from these plants, individuals may benefit from the protective effects against oxidative stress-related diseases<sup>31</sup>.

Plants rich in alkaloids continue to be valuable sources of bioactive compounds with diverse medicinal properties that contribute to human health and well-being. Many medicinal potentials of alkaloid compounds including antioxidant activity<sup>32</sup>, and antidiabetic activities have been reported.

The most significant amount of phytochemicals present in the bark of *Sterculia oblonga* stem is fatty acid/esters. This suggests that the plant could be a great source of health essential oil if properly extracted. The oils may help lower the adverse effects of diabetes and blood pressure<sup>34</sup>.

Plants containing hydrocarbons have shown potential therapeutic implications in various fields, primarily because of the existence of secondary metabolites that can exhibit medicinal properties. Some plants produce hydrocarbons that possess antimicrobial/anti-inflammatory properties<sup>35</sup> and anti-cancer potentials<sup>36</sup>

## CONCLUSION

The bark on the stem of *Sterculia oblonga* revealed the existence of twelve (12) different phytochemicals including organo-Sulphur compounds (2%), phenols (24%), alkaloids (2%), fatty acids/esters (52%) and hydrocarbons (20%). The presence of these compounds makes the plant a suitable choice for various therapeutic applications as far as herbal medicine is concerned.

Conflict of Interest: The authors declare no conflict of interest.

Authors' Contribution: All authors contributed to the preparation of the manuscript. All authors read and approved the final manuscript.

Funding: None.

## REFERENCIAS BIBLIOGRAFICAS

- [1] Lobo FC, Franco AR, Fernandes EM, Reis RL. An overview of the antimicrobial properties of lignocellulosic materials. *Molecules*. 2021 Mar 20;26(6):1749.
- [2] Ojo AO. An overview of lignocellulose and its biotechnological importance in high-value product production. *Fermentation*. 2023 Nov 20;9(11):990.
- [3] Cos P, Vlietinck AJ, Berghe DV, Maes L. Anti-infective potential of natural products: How to develop a stronger in vitro 'proof-of-concept'. *Journal of ethnopharmacology*. 2006 Jul 19;106(3):290-302.
- [4] Nassr-Allah AA, Aboul-Enein AM, Aboul-Enein KM, Lightfoot DA, Cocchetto A, El-Shemy HA. Anti-cancer and anti-oxidant activity of some Egyptian medicinal plants. *J Med Plants Res*. 2009 Oct 1;3(10):799-808.
- [5] Gil MI, Tomás-Barberán FA, Hess-Pierce B, Kader AA. Antioxidant capacities, phenolic compounds, carotenoids, and vitamin C contents of nectarine, peach, and plum cultivars from California. *Journal of agricultural and food chemistry*. 2002 Aug 14;50(17):4976-82.
- [6] Sasidharan S, Chen Y, Saravanan D, Sundram KM, Latha LY. Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African journal of traditional, complementary and alternative medicines*. 2011;8(1).
- [7] Bosh AH. and Louppe D. *Sterculia oblonga*.in: (Louppe D, Oteng-Ainoako AA and Brin M. *Plant Resource of Tropical Africa (PROTA)*, Wageningen, Netherlands. 2008.
- [8] Ezeonu CS, Ejikeme CM. Qualitative and quantitative determination of phytochemical contents of indigenous Nigerian softwoods. *New journal of Science*. 2016;2016(1):5601327.
- [9] Morah FN, Bisong EA. Chemical composition and testosterone level enhancing property of *Sterculia oblonga* leaf essential oil in male albino rats. *Public Health*. 2019;6(100024P16FM2019):100024P16FM2019.
- [10] Morah FN, Bisong EA. Chemical composition of *Sterculia oblonga* leaf petroleum spirit extract and its testosterone level enhancing property in male albino rats. *International Journal of Chemical and Biochemical Sciences*. 2021;19:89-93.
- [11] Shoaib S, Ansari MA, Ghazwani M, Hani U, Jamous YF, Alali Z, Wahab S, Ahmad W, Weir SA, Alomary MN, Yusuf N. Prospective epigenetic actions of organo-sulfur compounds against cancer: perspectives and molecular mechanisms. *Cancers*. 2023 Jan 23;15(3):697.
- [12] Abdelbasset WK, Elkholi SM, Ismail KA, AL-Ghamdi HS, Mironov S, Ridha HS, Maashi MS, Thangavelu L, Mahmudiono T, Mustafa YF. Mequinol-loaded carboxymethyl cellulose/chitosan electrospun wound dressing as a potential candidate to treat diabetic wounds. *Cellulose*. 2022 Sep;29(14):7863-81.
- [13] Almeida RS, Taccini MM, de Moura LF, Ceribelli UL, Brito JO, Gloria EM. Potential of pyroligneous extract of Eucalyptus wood as a preservative of cosmetic and sanitizing products. *Waste and Biomass Valorization*. 2019 May 1;10:1111-8.
- [14] Oramahi H, Yoshimura T, Rusmiyanto E, Kustiati K. Antifungal and antitermitic activities of vinegars from two biomass resources at different pyrolytic temperatures. *Journal of Applied Biological Sciences*. 2020 Apr 29;14(1):26-38.
- [15] Parmar VS, Gupta S, Bisht KS, Mukherjee S, Boll PM, Errington W. Synthesis and X-ray structure of 2-(3-methyl-2-butenyl)-3, 4, 5-trimethoxyphenol: a potent anti-invasive agent against solid tumours. *Acta Chemica Scandinavica*. 1996 Jun 1;50(6):558.
- [16] Verotta L, Dell'Agli M, Giolito A, Guerrini M, Cabalion P, Bosisio E. In vitro antiplasmodial activity of extracts of *Tristanopsis* species and identification of the active constituents: Ellagic acid and 3, 4, 5-trimethoxyphenyl-(6'-O-galloyl)-O-β-D-glucopyranoside. *Journal of natural products*. 2001 May 25;64(5):603-7.
- [17] Dardouk F, Arandi H, Makharzeh M. Extraction of Corn silk and its applications in healing and cosmetics'. Student project, 2019. An-Najah National University, Palestine.

- [18] Aparna V, Dileep KV, Mandal PK, Karthe P, Sadasivan C, Haridas M. Anti-inflammatory property of n-hexadecanoic acid: structural evidence and kinetic assessment. *Chemical biology & drug design*. 2012 Sep;80(3):434-9.
- [19] Ravi L, Krishnan K. Research article cytotoxic potential of N-hexadecanoic acid extracted from *Kigelia pinnata* leaves. *Asian J. Cell Biol*. 2017;12:20-7.
- [20] Keke CO, Nsofor WN, Kumabia FK, Iloabuchi GC, Ejiofor JC, Osuagwu OL. GCMS and FTIR analysis of ethanol and methanol leave extract of *Urena lobata* (Caesar weed) for bioactive phytochemical constituents. *Journal of Drug Delivery and Therapeutics*. 2023 Jan 15;13(1):99-115.
- [21] More K, Tayade S, Gawande P, Manik S, Shelke D. Antioxidant and antimicrobial potential of *Canavalia gladiata* (Jacq.) DC. leaves and seeds: GC-MS based metabolic profiling. *Indian Journal of Natural Products and Resources (IJNPR)[Formerly Natural Product Radiance (NPR)]*. 2022 Jul 26;13(2):163-9.
- [22] Jabeen N, Javaid A, Ahmed E. Antifungal activity and phytochemical profile of chloroform soluble fraction of *Datura metel* fruit. 2022. Pp 1085 – 1091.
- [23] Jacobson M, Ohinata K, Chambers DL, Jones WA, Fujimoto MS. Insect sex attractants. 13. Isolation, identification, and synthesis of sex pheromones of the male Mediterranean fruit fly. *Journal of medicinal chemistry*. 1973 Mar;16(3):248-51.
- [24] Banakar P, Jayaraj M. GC-MS analysis of bioactive compounds from ethanolic leaf extract of *Waltheria indica* Linn. and their pharmacological activities. *Int. J. Pharm. Sci. Res*. 2018 May 1;9(5):2005-10.
- [25] Banakar P, Jayaraj M. GC-MS analysis of bioactive compounds from ethanolic leaf extract of *Waltheria indica* Linn. and their pharmacological activities. *Int. J. Pharm. Sci. Res*. 2018 May 1;9(5):2005-10.
- [26] Sales-Campos H, Reis de Souza P, Crema Peghini B, Santana da Silva J, Ribeiro Cardoso C. An overview of the modulatory effects of oleic acid in health and disease. *Mini reviews in medicinal chemistry*. 2013 Feb 1;13(2):201-10.
- [27] Dilika F, Bremner PD, Meyer JJ. Antibacterial activity of linoleic and oleic acids isolated from *Helichrysum pedunculatum*: a plant used during circumcision rites. *Fitoterapia*. 2000 Aug 1;71(4):450-2.
- [28] Kamenetsky R, London Shafir I, Khassanov F, Kik C, Van Heusden AW, Vrielink-van Ginkel M, Burger-Meijer K, Auger J, Arnault I, Rabinowitch HD. Diversity in fertility potential and organo-sulphur compounds among garlics from Central Asia. *Biodiversity & Conservation*. 2005 Feb;14:281-95.
- [29] Petropoulos S, Di Gioia F, Ntatsi G. Vegetable organosulfur compounds and their health promoting effects. *Current pharmaceutical design*. 2017 May 1;23(19):2850-75.
- [30] Micová M, Bystrická J, Kovarovič J, Harangozo E, Lidiková A. Content of bioactive compounds and antioxidant activity in garlic (*Allium sativum* L.). *Acta agriculturae Slovenica*. 2018 Dec 12;111(3):581-95.
- [31] Kumar N, Goel N. Phenolic acids: Natural versatile molecules with promising therapeutic applications. *Biotechnology reports*. 2019 Dec 1;24:e00370.
- [32] Gan J, Feng Y, He Z, Li X, Zhang H. Correlations between antioxidant activity and alkaloids and phenols of maca (*Lepidium meyenii*). *Journal of Food Quality*. 2017;2017(1):3185945.
- [33] Tiong SH, Looi CY, Hazni H, Arya A, Paydar M, Wong WF, Cheah SC, Mustafa MR, Awang K. Antidiabetic and antioxidant properties of alkaloids from *Catharanthus roseus* (L.) G. Don. *Molecules*. 2013 Aug 15;18(8):9770-84.
- [34] Silva AP, da Costa WA, Salazar M, do Nascimento BP, Pires FC. Commercial and therapeutic potential of plant-based fatty acids. *IntechOpen*. 2018 Nov 5;5:73-90.
- [35] Perumal Samy R, Sethi G, TK Chow V, G Stiles B. Plant-based hydrocarbon esters from *Tragia involucrata* possess antimicrobial and anti-inflammatory activities. *Infectious Disorders-Drug Targets (Formerly Current Drug Targets-Infectious Disorders)*. 2013 Apr 1;13(2):141-53.
- [36] Al-Saraireh, Y. M., Youssef, A. M., Alsarayreh, A. Z., Al Hujran, T. A., Al-Sarayreh, S., Al-Shuneigat, J. M., & Alrawashdeh, H. M. (2021). Phytochemical and anti-cancer properties of *Euphorbia hierosolymitana* Boiss. crude extracts. *J. Pharm. Pharmacogn. Res*, 9(13-23).